arXiv:1411.1523v1 [math.AP] 6 Nov 2014

Some existence and nonexistence results for a Schrödinger-Poisson type system

Yutian Lei

Institute of Mathematics
School of Mathematical Sciences
Nanjing Normal University
Nanjing, 210023, China
Email:leiyutian@njnu.edu.cn

Abstract In this paper, we study the Schrödinger-Poisson system

$$\begin{cases} -\Delta u = \sqrt{p} u^{p-1} v, \quad u > 0 \quad in \quad R^n, \\ -\Delta v = \sqrt{p} u^p, \quad v > 0 \quad in \quad R^n \end{cases}$$

with $n \geq 3$ and $p > 1$. We investigate the existence and the nonexistence of positive classical solutions with the help of an integral system involving the Newton potential

$$\begin{cases} u(x) = c_1 \displaystyle\int_{R^n} \frac{u^{p-1}(y) v(y) dy}{|x-y|^{n-2}}, \quad u > 0 \quad in \quad R^n, \\ v(x) = c_2 \displaystyle\int_{R^n} \frac{u^p(y) dy}{|x-y|^{n-2}} \quad v > 0 \quad in \quad R^n. \end{cases}$$

First, the system has no solution when $p \leq \frac{n}{n-2}$. When $p > \frac{n}{n-2}$, the system has a singular solution on $R^n \setminus \{0\}$ with slow asymptotic rate $\frac{2}{p-1}$. When $p < \frac{n+2}{n-2}$, the system has no solution in $L^{\frac{n(p-1)}{2}}(R^n)$. In fact, if the system has solutions in $L^{\frac{n(p-1)}{2}}(R^n)$, then $p = \frac{n+2}{n-2}$ and all the positive classical solutions can be classified as $u(x) = v(x) = c(\frac{t}{t^2+|x-x^*|^2})^{\frac{n-2}{2}}$, where c, t are positive constants. When $p > \frac{n+2}{n-2}$, by the shooting method and the Pohozaev identity, we find another pair of radial solution (u, v) satisfying $u \equiv v$ and decaying with slow rate $\frac{2}{p-1}$.
Keywords: Schrödinger-Poisson equation, positive classical solution, existence and nonexistence, critical and noncritical conditions
MSC2010 35J10, 35J47, 35Q55, 45E10, 45G05

1 Introduction

Recently, many authors devoted to study the nonlocal stationary Schrödinger equation

$$-\Delta u = p u^{p-1} (|x|^{2-n} * u^p), \quad u > 0 \quad in \quad R^n, \tag{1.1}$$

where $n \geq 3$ and $p > 1$. In particular, Moroz and Van Schaftingen [25] studied the existence of the supersolutions and listed several sufficient conditions.

Equation (1.1) is defined in many places. One is the example 3.2.8 in the book [4]. A more general form is the Choquard type equation in the papers [14] and [24]. Another interesting work related to (1.1) is the paper [12] and the references therein. Equation (1.1) is also helpful in understanding the blowing up or the global existence and scattering of the solutions of the dynamic Hartree equation (cf. [18]), which arises in the study of boson stars and other physical phenomena, and also appears as a continuous-limit model for mesoscopic molecular structures in chemistry. Such an equation also arises in the Hartree-Fock theory of the nonlinear Schrödinger equations (cf. [23]). More related mathematical and physical background can be found in [2], [10], [26], [28] and the references therein.

Since (1.1) has a convolution term, it seems difficult to investigate the existence directly. Write

$$v(x) = \sqrt{p} \int_{R^n} \frac{u^p(y)dy}{|x-y|^{n-2}}.$$

Then $v > 0$ in R^n. Noting the relation between the Newton potential and the convolution properties of Dirac function, we see that

$$-\Delta v(x) = \sqrt{p}(-\Delta|x|^{2-n}) * u^p = \sqrt{p}\delta_x * u^p = \sqrt{p}u^p(x),$$

where δ_x is the Dirac mass at x. Thus, the positive solution of (1.1) must satisfy the following system

$$\begin{cases} -\Delta u = \sqrt{p}u^{p-1}v, \quad u > 0 \quad in \quad R^n, \\ -\Delta v = \sqrt{p}u^p, \quad v > 0 \quad in \quad R^n. \end{cases} \tag{1.2}$$

It is a simplified model of the Schrödinger-Piosson system (cf. [1], [11] and references therein).

Quittner and Souplet [27] studied positive solutions of another PDE system

$$\begin{cases} -\Delta u = v^p u^r, \quad u > 0 \quad in \quad R^n, \\ -\Delta v = v^s u^q, \quad v > 0 \quad in \quad R^n. \end{cases} \tag{1.3}$$

They proved the following results:

(R1) If $n \geq 3$, $p - s = q - r \geq 0$ and $0 \leq r, s \leq \frac{n}{n-2}$, then positive solutions u, v satisfy $u \equiv v$.

(R2) If $n \geq 3$, $p - s = q - r \geq 0$, then nonnegative solutions u, v satisfy $u \geq v$ or $v \geq u$.

In this paper, we study the existence of positive classical solutions to the Schrödinger-Piosson type system (1.2). Besides (R1) and (R2), we have further existence and nonexistence results and state them in three cases.

First we consider the subcritical case $p \in (1, 2^* - 1)$, where $2^* = \frac{2n}{n-2}$. We have the following nonexistence results.

Theorem 1.1. *(1)If $p \leq \frac{n}{n-2}$, then (1.2) has no positive solution.*

(2) If $2 \leq p < 2^* - 1$, then (1.2) has no positive solution in $L^{\frac{n(p-1)}{2}}(R^n)$.

Remark.

1. When $u \equiv v$, (1.2) is reduced to the Lane-Emden type equation

$$-\Delta u = \sqrt{p} u^p, \quad u > 0 \ in \ R^n. \tag{1.4}$$

Its nonexistence results similar to Theorem 1.1 can be found in [9]. For the system (1.2), we here introduce an integral system to investigate the nonexistence. By the same argument in [6] and [15], we know that the solution of (1.2) satisfies the following integral system involving Newton potentials

$$\begin{cases} u(x) = c_1 \displaystyle\int_{R^n} \frac{u^{p-1}(y)v(y)dy}{|x-y|^{n-2}}, \quad u > 0 \quad in \quad R^n, \\ v(x) = c_2 \displaystyle\int_{R^n} \frac{u^p(y)dy}{|x-y|^{n-2}} \quad v > 0 \quad in \quad R^n, \end{cases} \tag{1.5}$$

for some constants $c_1, c_2 > 0$. By this integral system we can verify the nonexistence of positive solutions when $p \leq \frac{n}{n-2}$. When $p > \frac{n}{n-2}$, then (1.2) always has singular solutions on $R^n \setminus \{0\}$ (cf. Theorem 4.1).

2. The integral system (1.5) is invariant after translation. Its $L^{\frac{n(p-1)}{2}}(R^n)$-solutions are radially symmetric about some point $x^* \in R^n$. In the non-critical case, Kelvin transformation breaks the translation invariant of (1.5). On the other hand, the $L^{\frac{n(p-1)}{2}}(R^n)$-solutions are still radially symmetric about the origin in the subcritical case. These facts lead to a contradiction and hence $L^{\frac{n(p-1)}{2}}(R^n)$-solution does not exist.

Next, we consider the critical case and classify the $L^{\frac{n(p-1)}{2}}(R^n)$-solutions.

Theorem 1.2. *Let u be a classical solution of (1.2). Then the following items are equivalent to each other*

(1) $u \in L^{\frac{n(p-1)}{2}}(R^n)$;
(2) u is bounded and decaying with the fast rate $n-2$;
(3) u belongs to the homogeneous Sobolev space $\mathcal{D}^{1,2}(R^n)$;
(4) $u \in L^{2^*}(R^n)$ and $p = 2^* - 1$;
(5) $u(x) = v(x) = c(\frac{t}{t^2+|x-x^*|^2})^{\frac{n-2}{2}}$ with constants $c, t > 0$.

In the special case $u \equiv v$, we recall the Lane-Emden equation (1.4). The classification of the solutions of this single equation has provided an important ingredient in the study of the conformal geometry, such as the prescribing scalar curvature problem and the extremal functions of the Sobolev inequalities. It was studied rather extensively (cf. [3], [8], [9], [21] and the reference therein). In particular, Chen and Li [5] proved that all the positive solutions of (1.4) with the critical exponent $p = 2^* - 1$ must be the form as

$$u(x) = c(\frac{t}{t^2 + |x-x^*|^2})^{\frac{n-2}{2}}$$

with constants $c, t > 0$. For the system (1.2), we expect to prove both $u \equiv v$ and $p = 2^* - 1$. Once these results are verified, we can use the result in [5] to classify the positive solutions of (1.2).

At last, we consider the supercritical case $p > 2^* - 1$. We use the shooting method and Pohozaev identity to prove the following result.

Theorem 1.3. *When $p > 2^* - 1$ and $p \geq 2$, we can find radial solutions u, v decaying with the slow rate $\frac{2}{p-1}$ when $|x| \to \infty$.*

Not all the solutions in the supercritical case are radially symmetric. In section 4 we introduce an example to show that some bounded solutions are neither radial nor decaying when $|x| \to \infty$.

Another integral system as (1.5) is the following Lane-Emden type equations

$$\begin{cases} u(x) = c_1 \displaystyle\int_{R^n} \frac{v^q(y)dy}{|x-y|^\lambda}, & u > 0 \quad in \quad R^n, \\ v(x) = c_2 \displaystyle\int_{R^n} \frac{u^p(y)dy}{|x-y|^\lambda}, & v > 0 \quad in \quad R^n. \end{cases} \tag{1.6}$$

It is essential in studying the extremal functions of the Hardy-Littlewood-Sobolev inequality (cf. [22])

$$\int_{R^n}\int_{R^n} \frac{f(x)g(y)}{|x-y|^\lambda}dxdy \leq C(n,s,\lambda)||f||_r||g||_s$$

with $0 < \lambda < n$, $1 < s, r < \infty$, $f \in L^r(R^n)$ and $g \in L^s(R^n)$, $\frac{1}{r} + \frac{1}{s} + \frac{\lambda}{n} = 2$.

Define $Tg(x) = \displaystyle\int_{R^n} \frac{g(y)}{|x-y|^{n-\alpha}}dy$ with $\alpha = n - \lambda$. The Hardy-Littlewood-Sobolev inequality becomes

$$||Tg||_p \leq C(n,s,\alpha)||g||_{\frac{np}{n+\alpha p}},$$

where $\frac{n}{n-\alpha} < p < \infty$, and $1 < s < n/\alpha$. This inequality will be used in this paper to research the radial symmetry and the integrability of the solutions of (1.5).

In the critical case, the classification results for the single equation of (1.6) can be found in [6] and [19]. In particular, the method of moving planes of integral forms was introduced in [6]. It has become a powerful tool to handle the qualitative properties including the existence and the nonexistence, the radial symmetry, and the priori estimates. Jin and Li [13] applied a regularity lifting lemma by the contraction maps to obtain the optimal integrability of positive regular solutions. Based on this result, [16] estimated the fast decay rates when $|x| \to \infty$.

2 Subcritical case

In this section, we always assume that $p < 2^* - 1$.

Theorem 2.1. *(Liouville theorem) If $p \le \frac{n}{n-2}$, then there does not exist any positive solution of (1.5).*

Proof. The idea in [15] can be used here.

If u, v are positive solutions, we can deduce a contradiction.

(1) When $p < \frac{n}{n-2}$, it follows $n - 2 - \frac{2}{p-1} < 0$.

Set $a_0 = n-2$, $a_{j+1} = (2p-1)a_j - 4$, $b_j = pa_j - 2$ for $j = 0, 1, 2, \cdots$.

By (1.5), we have

$$u(x) \ge c \int_{B_1(0)} \frac{u^{p-1}(y)v(y)dy}{|x-y|^{n-2}} \ge \frac{c}{|x|^{a_0}}, \tag{2.1}$$

$$v(x) \ge c \int_{B(x,|x|/2)} \frac{u^p(y)dy}{|x-y|^{n-2}} \ge \frac{c}{|x|^{b_0}},$$

$$u(x) \ge c \int_{B(x,|x|/2)} \frac{u^{p-1}(y)v(y)dy}{|x-y|^{n-2}} \ge \frac{c}{|x|^{a_1}},$$

$$\cdots \quad \cdots$$

By induction, we have $u(x) \ge \frac{c}{|x|^{a_j}}$ as long as $a_j > 0$. Noting $p > 1$ and $n - 2 - \frac{2}{p-1} < 0$, from

$$\begin{aligned} a_j &= a_0(2p-1)^j - 4[(2p-1)^{j-1} + (2p-1)^{j-2} + \cdots + (2p-1)^0] \\ &= (n - 2 - \tfrac{2}{p-1})(2p-1)^j + \tfrac{2}{p-1}, \end{aligned}$$

we can find some $j_0 > 0$ such that $a_{j_0} \le 0$. This leads to $u(x) = \infty$ and yields a contradiction.

(2) When $p = \frac{n}{n-2}$, from

$$u(x) \ge c \int_{B_R(0)} \frac{u^{p-1}(y)v(y)dy}{|x-y|^{n-2}} \ge \frac{c}{(R+|x|)^{n-2}} \int_{B_R(0)} u^{p-1}(y)v(y)dy,$$

we deduce

$$\begin{aligned} \int_{B_R(0)} u^p(x)dx &\ge \int_{B_R(0)} \frac{cdx}{(R+|x|)^n} (\int_{B_R(0)} u^{p-1}(y)v(y)dy)^p \\ &\ge c(\int_{B_R(0)} u^{p-1}(y)v(y)dy)^p. \end{aligned} \tag{2.2}$$

Here c is independent of R. Similarly, from

$$v(x) \ge \frac{c}{(R+|x|)^{n-2}} \int_{B_R(0)} u^p(y)dy, \tag{2.3}$$

we also deduce

$$\int_{B_R(0)} u^{p-1}(x)v(x)dx \geq \int_{B_R(0)} \frac{cu^{p-1}(x)dx}{(R+|x|)^{n-2}} \int_{B_R(0)} u^p(y)dy.$$

Using (2.1), (2.2) and noting $p = \frac{n}{n-2}$, we get

$$\int_{B_R(0)} u^{p-1}(x)v(x)dx \geq c(\int_{B_R(0)} u^{p-1}(y)v(y)dy)^p,$$

which implies $u^{p-1}v \in L^1(R^n)$ if we let $R \to \infty$.

Multiplying (2.3) by u^{p-1} and integrating on $A_R := B_R(0) \setminus B_{R/2}(0)$, we still have

$$\int_{A_R} u^{p-1}(x)v(x)dx \geq c(\int_{B_R(0)} u^{p-1}(y)v(y)dy)^p.$$

Letting $R \to \infty$ and noting $u^{p-1}v \in L^1(R^n)$, we obtain $\|u^{p-1}v\|_{L^1(R^n)} = 0$. It is impossible. □

By using the method of moving planes in integral forms, which was established by Chen-Li-Ou (cf. [6] and [7]), we prove a radial symmetry result.

Theorem 2.2. *Let $p \geq 2$. If $h \geq 0$ satisfies*

$$\int_{R^n} |y|^{-nh/2} u^{n(p-1)/2}(y)dy < \infty, \quad \int_{R^n} |y|^{-nh/2} v^{n(p-1)/2}(y)dy < \infty. \tag{2.4}$$

Then the positive continuous solutions of

$$\begin{cases} u(x) = c_1 \displaystyle\int_{R^n} \frac{u^{p-1}(y)v(y)dy}{|y|^h|x-y|^{n-2}}, \quad u > 0 \quad in \quad R^n, \\ v(x) = c_2 \displaystyle\int_{R^n} \frac{u^p(y)dy}{|y|^h|x-y|^{n-2}} \quad v > 0 \quad in \quad R^n, \end{cases} \tag{2.5}$$

are radially symmetric and decreasing about $x^* \in R^n$. Moreover, $x^* = 0$ as long as $h > 0$.

Proof. For some real number λ, define $\Sigma_\lambda := \{x = (x_1, \ldots, x_n); x_1 > \lambda\}$, $x^\lambda = (2\lambda - x_1, x_2, \ldots, x_n)$, $u_\lambda(x) = u(x^\lambda)$, $\Sigma_\lambda^u = \{x \in \Sigma_\lambda | u(x) \leq u_\lambda(x)\}$, $\Sigma_\lambda^v = \{x \in \Sigma_\lambda | v(x) \leq v_\lambda(x)\}$. It is not difficult to see that

$$\begin{aligned} & u_\lambda(x) - u(x) \\ &= c_1 \int_{\Sigma_\lambda} \left(\frac{1}{|x-y|^{n-2}} - \frac{1}{|x^\lambda - y|^{n-2}} \right) \frac{1}{|y^\lambda|^h} (v_\lambda u_\lambda^{p-1} - v u^{p-1}) dy \\ &- c_1 \int_{\Sigma_\lambda} \left(\frac{1}{|x-y|^{n-2}} - \frac{1}{|x^\lambda - y|^{n-2}} \right) \left(\frac{1}{|y|^h} - \frac{1}{|y^\lambda|^h} \right) v u^{p-1} dy. \end{aligned} \tag{2.6}$$

Since the second term of the right hand side is nonpositive, from the definition of Σ_λ^u and Σ_λ^v, it follows

$$\begin{aligned}
&u_\lambda(x) - u(x) \\
&\le c \int_{\Sigma_\lambda} \left(\frac{1}{|x-y|^{n-2}} - \frac{1}{|x^\lambda - y|^{n-2}} \right) \frac{1}{|y^\lambda|^h} (v_\lambda u_\lambda^{p-1} - v u^{p-1}) dy \\
&\le c \int_{\Sigma_\lambda^v} \frac{1}{|x-y|^{n-2}} \frac{1}{|y^\lambda|^h} u_\lambda^{p-1} (v_\lambda - v)(y) dy \\
&\quad + c \int_{\Sigma_\lambda^u} \frac{1}{|x-y|^{n-2}} \frac{1}{|y^\lambda|^h} v (u_\lambda^{p-1} - u^{p-1})(y) dy.
\end{aligned}$$

Using the Hardy-Littlewood-Sobolev inequality and the Hölder inequality, we have

$$\begin{aligned}
&\|u_\lambda - u\|_{L^s(\Sigma_\lambda^u)} \\
&\le C \| |y|^{-h} u_\lambda^{p-1} (v_\lambda - v) \|_{L^{\frac{ns}{n+2s}}(\Sigma_\lambda^v)} + C \| |y|^{-h} v u_\lambda^{p-2} (u_\lambda - u) \|_{L^{\frac{ns}{n+2s}}(\Sigma_\lambda^u)} \\
&\le C \| |y|^{-h} u_\lambda^{p-1} \|_{L^{\frac{n}{2}}(\Sigma_\lambda^v)} \| (v_\lambda - v) \|_{L^s(\Sigma_\lambda^v)} \\
&\quad + C \| |y|^{-h} u_\lambda^{p-2} v \|_{L^{\frac{n}{2}}(\Sigma_\lambda^u)} \| (u_\lambda - u) \|_{L^s(\Sigma_\lambda^u)}.
\end{aligned}$$

Similarly, we also obtain

$$\|v_\lambda - v\|_{L^s(\Sigma_\lambda^v)} \le C \| |y|^{-h} u_\lambda^{p-1} \|_{L^{\frac{n}{2}}(\Sigma_\lambda^u)} \| (u_\lambda - u) \|_{L^s(\Sigma_\lambda^u)}.$$

By (2.4), as $\lambda \to -\infty$,

$$C \| |y|^{-h} u_\lambda^{p-1} \|_{L^{\frac{n}{2}}(\Sigma_\lambda^v)} \le \frac{1}{4}, \quad \| |y|^{-h} u_\lambda^{p-2} v \|_{L^{\frac{n}{2}}(\Sigma_\lambda^u)} \le \frac{1}{4}.$$

Combining these results, we can see that Σ_λ^u and Σ_λ^v are empty set as long as λ is near $-\infty$.

Suppose that at $\lambda_0 < 0$, we have $u(x) \ge u_{\lambda_0}(x)$ and $v(x) \ge v_{\lambda_0}(x)$ but $u(x) \not\equiv u_{\lambda_0}(x)$ and $v(x) \not\equiv v_{\lambda_0}(x)$ on Σ_{λ_0}. By the same argument above, we can prove that there exists an $\epsilon > 0$, such that $u(x) \ge u_\lambda(x)$ and $v(x) \ge v_\lambda(x)$ on Σ_λ for all $\lambda \in [\lambda_0, \lambda_0 + \epsilon)$. Therefore, we can move plane $x_1 = \lambda$ to the right as long as

$$u(x) \ge u_\lambda(x) \quad \text{and} \quad v(x) \ge v_\lambda(x)$$

hold on Σ_λ. If the plane stops at $x_1 = \lambda_0$ for some $\lambda_0 < 0$, then $u(x)$ and $v(x)$ must be radially symmetric and decreasing about the plane $x_1 = \lambda_0$. Otherwise, we can move the plane all the way to $x_1 = 0$. Since the direction of x_1 can be chosen arbitrarily, we obtain that $u(x), v(x)$ are radially symmetric and decreasing about some $x^* \in R^n$.

If $h \neq 0$, we claim $x^* = 0$. Otherwise, we can find $\lambda_0 < 0$ such that $x_1 = \lambda_0$ is the stopped plane. From (2.6), we get

$$\begin{aligned} 0 &= u_{\lambda_0}(x) - u(x) \\ &= -c_1 \int_{\Sigma_{\lambda_0}} \left(\frac{1}{|x-y|^{n-2}} - \frac{1}{|x^{\lambda_0}-y|^{n-2}} \right) \left(\frac{1}{|y|^h} - \frac{1}{|y^{\lambda_0}|^h} \right) v u^{p-1} dy \\ &\neq 0. \end{aligned}$$

It is impossible. □

Theorem 2.3. *If $2 \leq p < 2^* - 1$, then there does not exist any positive solution of (1.5) in $L^{n(p-1)/2}(R^n)$.*

Proof. Step 1. Suppose u, v are the $L^{\frac{n(p-1)}{2}}(R^n)$-solutions of (1.5). According to Theorem 2.2 with $h = 0$, we see that they are radially symmetric about $x^* \in R^n$. Since (1.5) is invariant after translation, x^* can be chosen arbitrarily.

Step 2. Take the Kelvin transformation of u, v

$$\tilde{u}(x) = \frac{1}{|x|^{n-2}} u(\frac{x}{|x|^2}), \quad \tilde{u}(x) = \frac{1}{|x|^{n-2}} u(\frac{x}{|x|^2}).$$

By (1.5), we see that $\tilde{u}, \tilde{v}$ solve (2.5) with $h = n + 2 - p(n-2)$. In view of $p < 2^* - 1$, it follows $h > 0$. In addition, from $u, v \in L^{\frac{n(p-1)}{2}}(R^n)$, we see that (2.4) for $\tilde{u}, \tilde{v}$ is true. According to Theorem 2.2, $\tilde{u}, \tilde{v}$ are also radially symmetric but the center point x^* must be the origin. So the translation invariant is absent. By the same argument of Theorem 3 in [7], we can also deduce a contradiction. □

3 Critical case

Theorem 2.1 implies that

$$p > \frac{n}{n-2} \tag{3.1}$$

is the necessary condition of the existence of positive solutions for (1.2). Therefore, we can easily see that

$$u(x) = v(x) = c(\frac{t}{t^2 + |x - x^*|^2})^{\frac{n-2}{2}}$$

is a positive solution of (1.2) in $L^{n(p-1)/2}(R^n)$. However, it is nontrivial to show that all solutions of (1.2) in $L^{n(p-1)/2}(R^n)$ are the form above. In this section, we prove this conclusion.

Theorem 3.1. *If $u \in L^{\frac{n(p-1)}{2}}(R^n)$ solves (1.5), then*

(R1) $u, v \in L^s(R^n)$ for all $s > \frac{n}{n-2}$; $u, v \notin L^s(R^n)$ for all $s \leq \frac{n}{n-2}$;

(R2) u, v are bounded and

$$\lim_{|x|\to\infty} u(x) = \lim_{|x|\to\infty} v(x) = 0; \tag{3.2}$$

(R3) u, v are radially symmetric and decreasing about some point $x^* \in R^n$.

Proof. (1) By the Hardy-Littlewood-Sobolev inequality, from $u \in L^{\frac{n(p-1)}{2}}(R^n)$ we can deduce $v \in L^{\frac{n(p-1)}{2}}(R^n)$.

Write $w = u + v$. Then $w \in L^{\frac{n(p-1)}{2}}(R^n)$. From (1.5), it follows that w satisfies

$$w(x) = K(x) \int_{R^n} \frac{w^p(y)dy}{|x-y|^{n-2}}. \tag{3.3}$$

Here $K(x) > 0$ is upper bounded. Set $w_A = w$ as $|x| > A$ or $w > A$; $w = 0$ as $|x| \le A$ and $w \le A$. For $f \in L^s(R^n)$ with $s > \frac{n}{n-2}$, define

$$Tf(x) := K(x) \int_{R^n} \frac{w_A^{p-1}(y) f(y) dy}{|x-y|^{n-2}},$$
$$F(x) := K(x) \int_{R^n} \frac{(w-w_A)^p(y) dy}{|x-y|^{n-2}}.$$

Therefore, w solves the operator equation

$$f = Tf + F.$$

By the Hardy-Littlewood-Sobolev inequality, we get

$$\|Tf\|_s \le C \|w_A^{p-1} f\|_{\frac{ns}{n+2s}} \le C \|w_A\|^{p-1}_{\frac{n(p-1)}{2}} \|f\|_s,$$

and

$$\|F\|_s \le C \|w - w_A\|^p_{\frac{nps}{n+2s}} < \infty.$$

Take A suitably large such that $C\|w_A\|^{p-1}_{\frac{n(p-1)}{2}} < 1$. Thus, T is a contraction map from $L^s(R^n)$ to itself. In view of $\frac{n(p-1)}{2} > \frac{n}{n-2}$ (which is implied by (3.1)), T is also a contraction map from $L^{n(p-1)/2}(R^n)$ to itself. By the lifting lemma on the regularity (cf. Lemma 2.1 in [13]), we obtain $w \in L^s(R^n)$ for $s > \frac{n}{n-2}$. Thus, $u, v \in L^s(R^n)$ for $s > \frac{n}{n-2}$.

On the other hand, if $s \le \frac{n}{n-2}$, by (2.1) we have

$$\|u\|_s \ge c \int_R^\infty r^{n-s(n-2)} \frac{dr}{r} = \infty.$$

Similarly, we also deduce $v \notin L^s(R^n)$ for all $s \le \frac{n}{n-2}$. (R1) is proved.

(2) For $r > 0$, write

$$w(x) = K(x) \int_{B_r(x)} \frac{w^p(y)dy}{|x-y|^{n-2}} + K(x) \int_{R^n \setminus B_r(x)} \frac{w^p(y)dy}{|x-y|^{n-2}} := K_1 + K_2.$$

Clearly, for a suitably small $\epsilon > 0$, from (R1) we deduce

$$K_1 \le C\|w\|_{1/\epsilon}^p \Big(\int_0^d r^{n-\frac{n-2}{1-p\epsilon}} \frac{dr}{r}\Big)^{1-p\epsilon} \le C.$$

On the other hand, by virtue of (3.1) and (R1), we get $w \in L^p(R^n)$. Thus

$$K_2 \le Cr^{2-n}\|w\|_p^p \le C.$$

Combining the estimates of K_1 and K_2, we know that w is bounded. Thus, u, v are bounded.

Next, we show that w is decaying. Take $x_0 \in R^n$. By exchanging the order of the integral variables, we have

$$w(x_0) = (n-\alpha)K(x_0) \int_0^\infty \Big(\frac{\int_{B_t(x_0)} w^p(y)dy}{t^{n-\alpha}}\Big)\frac{dt}{t}.$$

Since $w \in L^\infty(R^n)$, $\forall \varepsilon > 0$, there exists $\delta \in (0, 1/2)$ such that

$$\int_0^\delta \Big[\frac{\int_{B_t(x_0)} w^p(z)dz}{t^{n-\alpha}}\Big]\frac{dt}{t} \le C\|w\|_\infty^p \int_0^\delta t^\alpha \frac{dt}{t} < \varepsilon. \tag{3.4}$$

As $|x - x_0| < \delta$,

$$\int_\delta^\infty \Big[\frac{\int_{B_t(x_0)} w^p(z)dz}{t^{n-\alpha}}\Big]\frac{dt}{t} \le \int_\delta^\infty \Big[\frac{\int_{B_{t+\delta}(x)} w^p(z)dz}{(t+\delta)^{n-\alpha}}\Big]\Big(\frac{t+\delta}{t}\Big)^{n-\alpha+1}\frac{d(t+\delta)}{t+\delta} \le Cw(x).$$

Combining this result with (3.4), we get

$$w(x_0) < C\varepsilon + Cw(x), \quad for \quad |x - x_0| < \delta.$$

Let $s = \frac{n(p-1)}{2}$, then

$$\begin{aligned} w^s(x_0) &= |B_\delta(x_0)|^{-1} \int_{B_\delta(x_0)} w^s(x_0)dx \\ &\le C\varepsilon^s + C|B_\delta(x_0)|^{-1}\|w\|^s_{L^s(B_\delta(x_0))}. \end{aligned} \tag{3.5}$$

Since $w \in L^s(R^n)$, $\lim_{|x_0|\to\infty} \|w\|_{L^s(B_\delta(x_0))} = 0$. Inserting this result into (3.5), we have

$$\lim_{|x_0|\to\infty} w^s(x_0) = 0.$$

This result means u, v converge to zero when $|x| \to \infty$. (R2) is proved.

(3) By the same proof of Theorem 2.2 with $h = 0$, we can see the conclusion (R3). □

Theorem 3.2. *If $u \in L^{\frac{n(p-1)}{2}}(R^n)$ solves (1.2) in the classical sense, then $u^p v \in L^1(R^n)$ and $u \in \mathcal{D}^{1,2}(R^n)$. Moreover,*

$$\int_{R^n} |\nabla u|^2 dx = \int_{R^n} |\nabla v|^2 dx = \sqrt{p}\int_{R^n} u^p v dx. \tag{3.6}$$

Proof. Step 1. By the Hölder inequality, from Theorem 3.1 (R1) and (3.1), we can deduce that $u^p v \in L^1(R^n)$.

Step 2. Take smooth function $\zeta(x)$ satisfying

$$\begin{cases} \zeta(x) = 1, & for\ |x| \le 1; \\ \zeta(x) \in [0,1], & for\ |x| \in [1,2]; \\ \zeta(x) = 0, & for\ |x| \ge 2. \end{cases}$$

Define the cut-off function

$$\zeta_R(x) = \zeta(\frac{x}{R}). \tag{3.7}$$

Multiplying the first equation of (1.2) by $u\zeta_R^2$ and integrating on $D := B_{3R}(0)$, we have

$$-\int_D \zeta_R^2 u \Delta u dx = \sqrt{p} \int_D u^p v \zeta_R^2 dx.$$

Integrating by parts, we obtain

$$\int_D |\nabla u|^2 \zeta_R^2 dx + 2\int_D u\zeta_R \nabla u \nabla \zeta_R dx = \sqrt{p} \int_D u^p v \zeta_R^2 dx. \tag{3.8}$$

Applying the Cauchy inequality, we get

$$|\int_D u\zeta_R \nabla u \nabla \zeta_R dx| \le \delta \int_D |\nabla u|^2 \zeta_R^2 dx + C\int_D u^2 |\nabla \zeta_R|^2 dx \tag{3.9}$$

for any $\delta \in (0, 1/2)$. According to Theorem 3.1 (R1), it follows $u, v \in L^{2^*}(R^n)$. By the Hölder inequality, we obtain

$$\int_D u^2 |\nabla \zeta_R|^2 dx \le (\int_D u^{2^*} dx)^{1-2/n} (\int_D |\nabla \zeta_R|^n dx)^{2/n} \le C. \tag{3.10}$$

Noting $u^p v \in L^1(R^n)$, from (3.8)-(3.10) we deduce

$$\int_D |\nabla u|^2 \zeta_R^2 dx \le C.$$

Letting $R \to \infty$ yields

$$\int_{R^n} |\nabla u|^2 dx < \infty.$$

Similarly, we also obtain

$$\int_{R^n} |\nabla v|^2 dx < \infty.$$

Combining the results above, we can see

$$\int_{R^n} (u^{2^*} + v^{2^*} + u^p v + |\nabla u|^2 + |\nabla v|^2) dx < \infty.$$

Therefore, we can find R_j such that

$$\lim_{R_j \to \infty} R_j \int_{\partial B_{R_j}} (u^{2^*} + v^{2^*} + u^p v + |\nabla u|^2 + |\nabla v|^2) ds = 0. \tag{3.11}$$

Step 3. Multiplying the first equation of (1.2) by u and integrating on D, we get

$$\int_D |\nabla u|^2 dx - \int_{\partial D} u\partial_\nu u ds = \sqrt{p}\int_D u^p v dx. \tag{3.12}$$

Here ν is the unit outward norm vector on ∂D. By the Hölder inequality, from (3.11) we deduce

$$|\int_{\partial D} u\partial_\nu u ds| \le CR^{\frac{n-1}{n}-1/2-1/2^*}(R\int_{\partial D}|\nabla u|^2 ds)^{1/2}(R\int_{\partial D}u^{2^*}ds)^{1/2^*} \to 0$$

when $R = R_j \to \infty$. Letting $R = R_j \to \infty$ in (3.12), we have $\|\nabla u\|_2^2 = \sqrt{p}\|u^p v\|_1$. Similarly, we can also obtain $\|\nabla v\|_2^2 = \sqrt{p}\|u^p v\|_1$. □

The following result shows that there does not exist $L^{\frac{n(p-1)}{2}}(R^n)$-solution if p is not equal to the critical exponent $2^* - 1$.

Theorem 3.3. *If a classical solution u belongs to $L^{\frac{n(p-1)}{2}}(R^n)$, then $p = 2^* - 1$, and hence $L^{n(p-1)/2}(R^n) = L^{2^*}(R^n)$.*

Proof. Write $B = B_R(0)$. Multiply (1.2) by $x\cdot\nabla u$ and $x\cdot\nabla v$, respectively. Integrating on B, we get

$$-\int_B (x\cdot\nabla u)\Delta u dx = \frac{1}{\sqrt{p}}\int_B v(x\cdot\nabla u^p)dx,$$

$$-\int_B (x\cdot\nabla v)\Delta v dx = \sqrt{p}\int_B u^p(x\cdot\nabla v)dx.$$

Integrating by parts yields

$$\begin{aligned}&-p\int_{\partial B}|x||\partial_\nu u|^2 ds + \frac{p}{2}\int_{\partial B}|x||\nabla u|^2 ds - \frac{n-2}{2}p\int_B |\nabla u|^2 dx\\ &= \sqrt{p}\int_B v(x\cdot\nabla u^p)dx,\end{aligned}$$

and

$$\begin{aligned}&-\int_{\partial B}|x||\partial_\nu v|^2 ds + \frac{1}{2}\int_{\partial B}|x||\nabla v|^2 ds - \frac{n-2}{2}\int_B |\nabla v|^2 dx\\ &= \sqrt{p}\int_B u^p(x\cdot\nabla v)dx.\end{aligned}$$

Adding two results together and integrating by parts again, we obtain

$$\begin{aligned}&-\int_{\partial B}|x|(p|\partial_\nu u|^2 + |\partial_\nu v|^2)ds + \frac{1}{2}\int_{\partial B}|x|(p|\nabla u|^2 + |\nabla v|^2)ds\\ &-\frac{n-2}{2}\int_B (p|\nabla u|^2 + |\nabla v|^2)dx\\ &= \sqrt{p}\int_B x\cdot\nabla(u^p v)dx = \sqrt{p}\int_{\partial B}|x|u^p v ds - n\sqrt{p}\int_B u^p v dx.\end{aligned}$$

Letting $R = R_j \to \infty$ and using (3.11), we have

$$\frac{n-2}{2}\int_B (p|\nabla u|^2 + |\nabla v|^2)dx = n\sqrt{p}\int_B u^p v dx.$$

By (3.6) we see $p = 2^* - 1$ finally. □

Theorem 3.4. *If a classical solution u of (1.2) belongs to $L^{\frac{n(p-1)}{2}}(R^n)$, then*

$$u(x) = v(x) = c(\frac{t}{t^2 + |x - x^*|^2})^{\frac{n-2}{2}}. \tag{3.13}$$

with some constant $c = c(n)$ and for some $t > 0$.

Proof. Step 1. We claim $u \equiv v$.

Let $W = u - v$. By Theorems 3.1 and 3.2, we see

$$\int_{R^n} (|W|^{2^*} + |\nabla W|^2)dx < \infty.$$

Thus, when $R = R_j \to \infty$,

$$R\int_{\partial B_R(0)} (|W|^{2^*} + |\partial_\nu W|^2)ds \to 0.$$

From (1.2), it follows $\Delta W = \sqrt{p}u^{p-1}W$. Therefore,

$$\int_B |\nabla W|^2 dx + \sqrt{p}\int_B u^{p-1}W^2 dx = \int_{\partial B} W\partial_\nu W ds. \tag{3.14}$$

Here $B = B_R(0)$. By the Hölder inequality, as $R = R_j \to \infty$,

$$\begin{aligned} &|\int_{\partial B} W\partial_\nu W ds| \\ &\le C(R\int_{\partial B} |W|^{2^*} ds)^{1/2^*} (R\int_{\partial B} |\partial_\nu W|^2 ds)^{1/2} R^{(n-1)(1/2-1/2^*)-1/2-1/2^*} \\ &\to 0. \end{aligned}$$

Inserting this into (3.14) with $R = R_j \to \infty$, we get

$$\int_{R^n} (|\nabla W|^2 + \sqrt{p}u^{p-1}W^2)dx = 0,$$

which implies $u \equiv v$.

Step 2. By virtue of $u \equiv v$ and Theorem 3.3, (1.2) is reduced to the single equation

$$-\Delta u = \sqrt{p}u^{2^*-1}, \quad u > 0 \text{ in } R^n.$$

According to the classification results in [5], the positive solutions must be the form as (3.13) in the critical case. □

The argument above implies $u \in L^{\frac{n(p-1)}{2}}(R^n)$ is equivalent to (3.13).

At last, we complete the proof of Theorem 1.2.

Theorem 3.5. *If the classical solution u of (1.2) belongs to $L^{\frac{n(p-1)}{2}}(R^n)$, if and only if one of the following items holds*

(C1) $u \in L^{2^*}(R^n)$ and $p = 2^* - 1$;

(C2) $u \in L^{\infty}(R^n)$ and $u(x) = O(|x|^{2-n})$ as $|x| \to \infty$;

(C3) $u \in \mathcal{D}^{1,2}(R^n)$.

Proof. The necessity can be seen in Theorems 3.1-3.4. Next, we state the sufficiency.

Both (C1) and (C2) can lead easily to $u \in L^{\frac{n(p-1)}{2}}(R^n)$. So, $u \in L^{\frac{n(p-1)}{2}}(R^n)$ is equivalent to (C1) and (C2).

Finally, (C3) implies (C1). In fact, by the Sobolev inequality we get $u \in L^{2^*}(R^n)$. On the other hand, using (3.8)-(3.10) we can deduce $u^p v \in L^1(R^n)$ from $u \in \mathcal{D}^{1,2}(R^n)$. Thus, (3.11) holds, and the proof of Theorem 3.3 still works. This shows $p = 2^* - 1$. □

Remark. $u(x) = O(|x|^{2-n})$ as $|x| \to \infty$ in condition (C2) implies $0 < c_1 \le u(x)|x|^{n-2} \le c_2$ as $|x| \to \infty$. It is led to by (2.1).

4 Supercritical case

If (3.1) holds, then (1.2) has a singular solution.

Theorem 4.1. *Let $u(x) = \frac{c_1}{|x|^{t_1}}$ and $v(x) = \frac{c_2}{|x|^{t_2}}$ solve (1.2) on $R^n \setminus \{0\}$, where c_1, c_2, t_1, t_2 are positive constants. Then such singular solutions must be*

$$u(x) = v(x) = \frac{c}{|x|^{\frac{2}{p-1}}}$$

with $c = [\frac{2n}{\sqrt{p}(p-1)} - \frac{4\sqrt{p}}{(p-1)^2}]^{\frac{1}{p-1}}$.

Proof. Write $U(r) = U(|x|) = u(x)$ and $V(r) = V(|x|) = v(x)$. Thus,

$$-U'' - \tfrac{n-1}{r}U' = \tfrac{c_1 t_1}{r^{t_1+2}}(n - t_1 - 2),$$

$$-V'' - \tfrac{n-1}{r}V' = \tfrac{c_2 t_2}{r^{t_2+2}}(n - t_2 - 2).$$

Since u, v solve (1.2), it is easy to get

$$\begin{cases} t_1 + 2 = t_1(p-1) + t_2, \\ t_2 + 2 = t_1 p, \end{cases}$$

and

$$\begin{cases} c_1 t_1 (n - t_1 - 2) = \sqrt{p} c_1^{p-1} c_2, \\ c_2 t_2 (n - t_2 - 2) = \sqrt{p} c_1^{p} \end{cases}$$

as long as $n - 2 > \max\{t_1, t_2\}$. Therefore, $t_1 = t_2 = \frac{2}{p-1}$ and $c_1 = c_2 = [\frac{2n}{\sqrt{p}(p-1)} - \frac{4\sqrt{p}}{(p-1)^2}]^{\frac{1}{p-1}}$. In view of (3.1), the condition $n - 2 > \max\{t_1, t_2\}$ is true obviously. □

Remark. The fact $u \equiv v$ is natural by an analogous argument in Step 4 of the proof of Theorem 4.4.

Next, we search for bounded solutions.

An example is the pair of cylinder-shaped solutions (u^*, v^*) (cf. [7]). Let

$$u_*(x) = c(\frac{t}{t^2 + |x - x^*|^2})^{\frac{n-2}{2}}. \tag{4.1}$$

According to Theorem 1.2, u_* solves (1.2) in the whole space R^n in the critical case $p = \frac{n+2}{n-2}$. Thus, it is not difficult to see that $u^*(x, x_{n+1}) = u_*(x)$ and $v^*(x_0, x) = u_*(x)$ still solves (1.2) in R^{n+1}. In view of $p > \frac{n+3}{n-1}$, the problem (1.2) which u^* satisfies in R^{n+1} is equipped with the supercritical exponent. Clearly, this pair of positive solution (u^*, v^*) is neither radial nor decaying when $|x| \to \infty$. We also see $u^* \neq v^*$ because the generating lines of the cylinders are different.

At last, we search for a radial bounded solution decaying with the slow rate $\frac{2}{p-1}$.

In order to find the existence of entire solutions in R^n, we need the following nonexistence result on a bounded domain. It can be verified by the Pohozaev identity.

Theorem 4.2. *Let $D \subset R^n$ be a ball centered at the origin. If*

$$p \geq 2^* - 1, \tag{4.2}$$

then the following boundary value problem has no positive radial solution in $C^2(D) \cap C^1(\bar{D})$

$$-\Delta u = \sqrt{p} u^{p-1} v, \quad in \quad D, \tag{4.3}$$

$$-\Delta v = \sqrt{p} u^{p}, \quad in \quad D, \tag{4.4}$$

$$u = v = 0 \quad on \quad \partial D. \tag{4.5}$$

Proof. Suppose that u, v are positive radial solutions, we will deduce a contradiction.

Multiply (4.3) and (4.4) by u and v, respectively. Integrating on D and using (4.5), we have

$$\int_D |\nabla u|^2 dx = \int_D |\nabla v|^2 dx = \sqrt{p} \int_D u^p v dx. \tag{4.6}$$

Since u has the radial structure, $|\nabla u|^2 = |\partial_\nu u|^2$ on ∂D. Here ν is the unit outward norm vector on ∂D.

Multiplying (4.3) by $(x \cdot \nabla u)$ and integrating on D, we get

$$-\int_{\partial D} |x||\partial_\nu u|^2 ds + \int_D |\nabla u|^2 dx + \frac{1}{2}\int_D x \cdot \nabla(|\nabla u|^2) dx = \frac{1}{\sqrt{p}}\int_D v(x \cdot \nabla u^p) dx.$$

Integrating by parts and noting (4.5), we obtain

$$\frac{1}{2}\int_{\partial D} |x||\partial_\nu u|^2 ds + \frac{n-2}{2}\int_D |\nabla u|^2 dx = \frac{n}{\sqrt{p}}\int_D u^p v dx + \frac{1}{\sqrt{p}}\int_D u^p (x \cdot \nabla v) dx.$$

Similarly, from (4.4) we also deduce that

$$-\frac{1}{2}\int_{\partial D} |x||\partial_\nu v|^2 ds - \frac{n-2}{2}\int_D |\nabla v|^2 dx = \sqrt{p}\int_D u^p (x \cdot \nabla v) dx.$$

Combining two results above with (4.6) yields

$$-\frac{1}{2}\int_{\partial D} |x|(|\partial_\nu v|^2 + \frac{1}{p}|\partial_\nu v|^2) ds = \frac{n-2}{2}(\sqrt{p} + \frac{1}{\sqrt{p}})\int_D u^p v dx - \frac{n}{\sqrt{p}}\int_D u^p v dx.$$

Therefore,

$$\frac{n-2}{2}(\sqrt{p} + \frac{1}{\sqrt{p}}) - \frac{n}{\sqrt{p}} < 0,$$

which contradicts with (4.2). □

Based on the Liouville type result above, we can search for positive solutions of (1.2) with radial structures. Let u, v be radially symmetric about $x^* \in R^n$. We can write

$$U(r) = U(|x - x^*|) = u(x - x^*),$$

$$V(r) = V(|x - x^*|) = v(x - x^*).$$

Theorem 4.3. *Let $p \geq \max\{2, 2^* - 1\}$. Then the following ODE system*

$$\begin{cases} -(U'' + \frac{n-1}{r}U') = \sqrt{p}U^{p-1}V, \quad -(V'' + \frac{n-1}{r}V') = \sqrt{p}U^p, \quad r > 0 \\ U'(0) = V'(0) = 0, \quad U(0) = 1, \quad V(0) = a, \end{cases} \tag{4.7}$$

has entire solutions for some positive constant a.

Proof. Here we use the shooting method.

Step 1. By the standard contraction argument, by $p \geq 2$ we can see the local existence. We denote the solutions by $u_a(r), v_a(r)$.

Step 2. We claim that either (4.7) has entire solutions for all $a > 1$, or for some $a^* > 1$, there exists $R \in (0, 1]$ such that $u_{a^*}(r), v_{a^*}(r) > 0$ for $r \in [0, R)$ and $u_{a^*}(R) = 0$.

In fact, integrating (4.7) twice yields

$$v_a(r) = v_a(0) - \sqrt{p}\int_0^r \tau^{1-n}\int_0^\tau s^{n-1}u_a^p(s) ds d\tau,$$

and

$$u_a(r) = u_a(0) - \sqrt{p}\int_0^r \tau^{1-n}\int_0^\tau s^{n-1}u_a^{p-1}(s)v_a(s)dsd\tau.$$

Thus,

$$v_a(r)-u_a(r) = (a-1)+\sqrt{p}\int_0^r \tau^{1-n}\int_0^\tau s^{n-1}u_a^{p-1}(s)(v_a(s)-u(s))dsd\tau. \quad (4.8)$$

Let $a > 1$. So we can find $\delta > 0$ such that $v_a(r) > u_a(r)$ for $r \in [0, \delta)$ by the continuity of u_a, v_a.

We claim $v_a(r) > u_a(r)$ for all $r \geq 0$. Otherwise, there exists $r_0 \geq \delta$ such that $v(r_0) = u(r_0)$ and $v(r) > u(r)$ as $r \in [0, r_0)$. From (4.8) with $r = r_0$ we can deduce a contradiction easily.

Therefore, if $u_a(r) > 0$ for all $r \geq 0$, then (4.7) has entire solutions and the proof is complete. Otherwise, we can find $R > 0$ such that $u_a(r), v_a(r) > 0$ for $r \in (0, R)$ and $u_a(R) = 0$. We denote the a in this state by a^*.

Step 3. We claim that for $a < \varepsilon_0 = \frac{1}{n2^{1+p}\sqrt{p}}$, there exists $R \in (0, 1]$, such that $u_a(r), v_a(r) > 0$ for $r \in [0, R)$ and $v_a(R) = 0$.

In fact, from (4.7) we obtain $u(r) > v(r)$ for $r \geq 0$ and $u_a', v_a' < 0$ for $r > 0$. Thus, $v_a(r) \leq a < \varepsilon_0$ and $u_a(r) \leq u_a(0) = 1$. Thus,

$$u_a(r) = u_a(0) - \sqrt{p}\int_0^r \tau^{1-n}\int_0^\tau s^{n-1}u_a^{p-1}(s)v_a(s)dsd\tau \geq 1 - \frac{\varepsilon_0\sqrt{p}r^2}{2n} \geq \frac{1}{2},$$

for $r \in (0, 1)$. Therefore,

$$v_a(r) = v_a(0) - \sqrt{p}\int_0^r \tau^{1-n}\int_0^\tau s^{n-1}u_a^{p}(s)dsd\tau < \varepsilon_0 - \frac{\sqrt{p}}{2^p}\frac{r^2}{2n}.$$

This proves that for $a < \varepsilon_0$, we can find $R \in (0, 1]$ such that $u_a(r), v_a(r) > 0$ for $r \in (0, R)$ and $v_a(R) = 0$.

Step 4. Let $\underline{a} = \sup \underline{S}$, where

$$\begin{aligned}\underline{S} := \{\varepsilon;\quad & \text{when } a \in (0, \varepsilon), \quad \exists R_a > 0, \text{ such that} \\ & u_a(r) > 0, v_a(r) \geq 0, \text{ for } r \in [0, R_a], v_a(R_a) = 0\}.\end{aligned}$$

Clearly, $\underline{S} \neq \emptyset$ by virtue of $\varepsilon_0 \in \underline{S}$. From Step 2, it follows $\varepsilon \leq a^*$ for $\varepsilon \in \underline{S}$. Namely, $\underline{S}$ is upper bounded, and hence we see the existence of $\underline{a}$.

Step 5. Write $\bar{u}(r) = u_{\underline{a}}(r)$ and $\bar{v}(r) = v_{\underline{a}}(r)$. We claim that $\bar{u}(r), \bar{v}(r) > 0$ for $r \in [0, \infty)$, and hence they are entire positive solutions of (4.7).

Otherwise, there exists $\bar{R} > 0$ such that $\bar{u}(r), \bar{v}(r) > 0$ for $r \in (0, \bar{R})$ and one of the following consequences holds:

$$(1)\ \ \bar{u}(\bar{R}) = 0, \bar{v}(\bar{R}) > 0;$$

$$(2)\ \ \bar{v}(\bar{R}) = 0, \bar{u}(\bar{R}) > 0;$$

$$(3)\ \ \bar{u}(\bar{R}) = 0, \bar{v}(\bar{R}) = 0.$$

We deduce the contradictions from three consequences above.

(1) By C^1-continuous dependence of u_a, v_a in a, and the fact $\bar{u}'(\bar{R}) < 0$, we see that for all $|a - \underline{a}|$ small, there exists $R_a > 0$ such that

$$\begin{aligned} &\bar{u}(r), \bar{v}(r) > 0, \quad for\ r \in (0, R_a); \\ &\bar{u}(R_a) = 0, \quad \bar{v}(R_a) > 0. \end{aligned}$$

This contradicts with the definition of $\underline{a}$.

(2) Similarly, for $|a - \underline{a}|$ small, there exists $R_a > 0$ such that

$$\begin{aligned} &\bar{u}(r), \bar{v}(r) > 0, \quad for\ r \in (0, R_a); \\ &\bar{u}(R_a) > 0, \quad \bar{v}(R_a) = 0. \end{aligned}$$

This implies that $\underline{a} + \delta \in \underline{S}$ for some $\delta > 0$, which contradicts with the definition of $\underline{a}$.

(3) The consequence implies that $u(x) = \bar{u}(|x|)$ and $v(x) = \bar{v}(|x|)$ are solutions of the system

$$\begin{cases} -\Delta u = \sqrt{p} u^{p-1} v, \quad -\Delta v = \sqrt{p} u^p, \ in\ B_R, \\ u, v > 0\ in\ B_R, \quad u = v = 0\ on\ \partial B_R. \end{cases} \tag{4.9}$$

It is impossible by Theorem 4.2.

All the contradictions show that our claim is true. Thus, the entire positive solutions exist. □

Theorem 4.4. *The entire solutions obtained in Theorem 4.3 satisfy*

$$\lim_{|x| \to \infty} u(x) = \lim_{|x| \to \infty} v(x) = 0 \tag{4.10}$$

and $u \equiv v$. Moreover, either

$$p = 2^* - 1 \quad and \quad u = u_*,$$

or

$$p > 2^* - 1 \quad and \quad c_1 \le u(x)|x|^{\frac{2}{p-1}} \le c_2$$

when $|x| \to \infty$, where c_1, c_2 are positive constants.

Proof. Step 1. We claim $\lim_{r \to \infty} \bar{u}(r) = 0$.

Eq. (4.7) implies $\bar{u}' < 0$ and $\bar{v}' < 0$ for $r > 0$. So $\bar{u}$ and $\bar{v}$ are decreasing positive solutions, and $\lim_{r \to \infty} \bar{u}(r)$, $\lim_{r \to \infty} \bar{v}(r)$ exist.

If there exists $c > 0$ such that $\bar{u}(r) \ge c$ for $r > 0$, then (4.7) shows that $\bar{v}$ satisfies

$$\bar{v}'' + \frac{n-1}{r} \bar{v}' \le -\sqrt{p} c^p.$$

Integrating twice yields

$$\bar{v}(r) \le \bar{v}(0) - \frac{\sqrt{p} c^p r^2}{2n}$$

for $r > 0$. It is impossible since $\bar{v}$ is an entire positive solution. This shows that $\bar{u} \to 0$ when $r \to \infty$.

Step 2. Furthermore, we claim $u(x) \le c_2|x|^{\frac{-2}{p-1}}$ when $|x| \to \infty$.

In fact, since $u(x), v(x)$ are the positive entire solutions of (1.2), they satisfy (1.5) according to the Remark in section 1. Moreover, $u(x), v(x)$ are radially symmetric and decreasing, we can deduce

$$v(x) \ge cu^p(x) \int_0^{|x|} r^2 \frac{dr}{r} \ge c|x|^2 u^p(x),$$

and hence

$$u(x) \ge cu^{p-1}(x)v(x) \int_0^{|x|} r^2 \frac{dr}{r} \ge c|x|^2 u^{p-1}(x)v(x) \ge c|x|^4 u^{2p-1}(x).$$

This implies our claim.

Step 3. We claim $\lim_{|x|\to\infty} v(x) = 0$.

Clearly, for large $R > 0$, we obtain from (1.5) that

$$\begin{aligned} v(x) &= c_2\Big(\int_{B_R(0)} \frac{u^p(y)dy}{|x-y|^{n-2}} + \int_{B_{|x|/2}(x)} \frac{u^p(y)dy}{|x-y|^{n-2}} \\ &\quad + \int_{B_R^c(0)\setminus B_{|x|/2}(x)} \frac{u^p(y)dy}{|x-y|^{n-2}}\Big) \\ &:= c_2(I_1 + I_2 + I_3). \end{aligned}$$

First, u is bounded. So, as $|x| \to \infty$,

$$I_1 \le C \int_{B_R(0)} \frac{dy}{|x-y|^{n-2}} \le C|x|^{2-n} \to 0.$$

Next, for large $|x|$, Step 2 implies $u(y) \le c|x|^{\frac{-2}{p-1}}$ as $y \in B_{|x|/2}(x)$. Therefore,

$$I_2 \le c|x|^{\frac{-2p}{p-1}} \int_{B_{|x|/2}(x)} \frac{dy}{|x-y|^{n-2}} \le C|x|^{2-\frac{2p}{p-1}} \to 0.$$

Finally, using the Young inequality and letting $R \to \infty$, we have

$$I_3 \le C \int_{B_R^c(0)\setminus B_{|x|/2}(x)} \frac{dy}{|x-y|^{n-2}|y|^{\frac{2p}{p-1}}} \le C \int_R^\infty r^{2-\frac{2p}{p-1}} \frac{dr}{r} \to 0.$$

Combining the estimates of I_1, I_2 and I_3, we get

$$\lim_{|x|\to\infty} v(x) = 0.$$

This result, together with Step 1, implies (4.10).

Step 4. We claim $u \equiv v$. The argument in Step 1 of the proof of Theorem 3.4 does not work, since the boundary integral is difficult to handle. We use

the idea in [17] to prove this uniqueness. This idea is universally valid, which implies the uniqueness as long as (4.10) holds. In addition, this idea is also independent whether the value of p is critical or not.

For any $r_0 \geq 0$, we prove $U(r_0) = V(r_0)$. Otherwise, either

$$U(r_0) < V(r_0), \tag{4.11}$$

or

$$U(r_0) > V(r_0). \tag{4.12}$$

If (4.11) holds, by the continuity of U and V, we can find $R > 0$ such that

$$U(r) < V(r) \quad as \quad r \in [r_0, R). \tag{4.13}$$

Set $R_0 = \sup\{R; (4.13) \text{ is true}\}$. Thus, when $R_0 < \infty$,

$$U(R_0) = V(R_0). \tag{4.14}$$

In view of (4.10), this conclusion still holds even if $R_0 = \infty$.

By (1.2), we see that for $r > 0$,

$$-(r^{n-1}U')' = \sqrt{p}r^{n-1}U^{p-1}V, \quad -(r^{n-1}V')' = \sqrt{p}r^{n-1}U^p.$$

Integrating twice, we get

$$U(R_0) = U(r_0) - \sqrt{p}\int_{r_0}^{R_0} r^{1-n}\int_0^r s^{n-1}U^{p-1}(s)V(s)dsdr,$$

$$V(R_0) = V(r_0) - \sqrt{p}\int_{r_0}^{R_0} r^{1-n}\int_0^r s^{n-1}U^{p}(s)dsdr.$$

Using (4.11), (4.13) and (4.14), we obtain

$$0 > U(r_0) - V(r_0) = \sqrt{p}\int_{r_0}^{R_0} r^{1-n}\int_0^r s^{n-1}U^{p-1}(s)(V(s) - U(s))dsdr \geq 0,$$

which is impossible. So (4.11) is not true. Similarly, we also see that (4.12) is not true. Since r_0 is arbitrary, we know $u \equiv v$.

Step 5. By virtue of $u \equiv v$, (1.2) is reduced to the single equation

$$-\Delta u = \sqrt{p}u^p, \quad u > 0 \ in \ R^n$$

with (4.2). According to the results in [20] and [21], we know that u either decays fast

$$c_1 \leq u(x)|x|^{n-2} \leq c_2,$$

or decays slowly

$$c_1 \leq u(x)|x|^{\frac{2}{p-1}} \leq c_2,$$

when $|x| \to \infty$. Here c_1, c_2 are positive constants.

If u decays fast, by Theorem 1.2 we know $p = 2^* - 1$ and $u \equiv u_*$. Here u_* is the radial function in (4.1).

If u decays slowly, we claim $p > 2^* - 1$. Otherwise, from (4.2) we have $p = 2^* - 1$. According to the classification result in [5], $u \equiv u_*$. This contradicts the slow decay rate. □

References

[1] A. Ambrosetti, *On Schrödinger-Poisson systems*, Milan J. Math., **76** (2008), 257–274.

[2] N. Ben Abdallah, F. Mehats, *On a Vlasov-Schrödinger-Poisson model*, Comm. Partial Differential Equations, **29** (2004), 173–206.

[3] L. Caffarelli, B. Gidas, J. Spruck, *Asymptotic symmetry and local behavior of semilinear elliptic equations with critical Sobolev growth*, Comm. Pure Appl. Math. **42** (1989), 271–297.

[4] T. Cazenave, *Semilinear schrödinger equations*, Courant Lecture Notes in Mathematics, 10. New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 2003.

[5] W. Chen, C. Li, *Classification of solutions of some nonlinear elliptic equations*, Duke Math. J., **63** (1991), 615–622.

[6] W. Chen, C. Li, B. Ou, *Classification of solutions for an integral equation*, Comm. Pure Appl. Math., **59** (2006), 330–343.

[7] W. Chen, C. Li, B. Ou, *Qualitative properties of solutions for an integral equation*, Discrete Contin. Dyn. Syst., **12** (2005), 347–354.

[8] B. Gidas, W.-M. Ni, L. Nirenberg, *Symmetry of positive solutions of nonlinear elliptic equations in R^n* (collected in the book *Mathematical Analysis and Applications*, which is vol. 7a of the book series *Advances in Mathematics. Supplementary Studies*, Academic Press, New York, 1981.)

[9] B. Gidas, J. Spruck, *Global and local behavior of positive solutions of nonlinear elliptic equations*, Comm. Pure Appl. Math., **34** (1981), 525–598.

[10] J. Ginibre, G. Velo, *Long range scattering and modified wave operators for some Hartree type equations*, Rev. Math. Phys., **12** (2000), 361–429.

[11] E. Hebey, J. Wei, *Schrödinger-Poisson systems in the 3-sphere*, Calc. Var. Partial Differential Equations, **47** (2013), 25–54.

[12] L. Jeanjean, T. Luo, *Sharp nonexistence results of prescribed L^2-norm solutions for some class of Schrödinger-Poisson and quasi-linear equations*, Z. Angew. Math. Phys., **64** (2013), 937–954.

[13] C. Jin, C. Li, *Qualitative analysis of some systems of integral equations*, Calc. Var. Partial Differential Equations, **26** (2006), 447–457.

[14] Y. Lei, *On the regularity of positive solutions of a class of Choquard type equations*, Math. Z., **273** (2013), 883–905.

[15] Y. Lei, *Qualitative analysis for the static Hartree-type equations*, SIAM J. Math. Anal., **45** (2013), 388–406.

[16] Y. Lei, C. Li, C. Ma, *Asymptotic radial symmetry and growth estimates of positive solutions to weighted Hardy-Littlewood-Sobolev system*, Calc. Var. Partial Differential Equations, **45** (2012), 43–61.

[17] C. Li, L. Ma, *Uniqueness of positive bound states to Schrödinger systems with critical exponents*, SIAM J. Math. Anal., **40** (2008), 1049–1057.

[18] D. Li, C. Miao, X. Zhang, *The focusing energy-critical Hartree equation*, J. Differential Equations, **246** (2009), 1139–1163.

[19] Y.-Y. Li, *Remark on some conformally invariant integral equations: the method of moving spheres*, J. Eur. Math. Soc., **6** (2004), 153–180.

[20] Y. Li, *Asymptotic behavior of positive solutions of equation* $\Delta u + K(x)u^p = 0$ *in* R^n, J. Differential Equations, **95** (1992), 304–330.

[21] Y. Li, W.-M. Ni, *On conformal scalar curvature equations in* R^n, Duke Math. J., **57** (1988), 895–924.

[22] E. Lieb, *Sharp constants in the Hardy-Littlewood-Sobolev and related inequalities*, Ann. of Math., **118** (1983), 349–374.

[23] E. Lieb, B. Simon, *The Hartree-Fock theory for Coulomb systems*, Comm. Math. Phys., **53** (1977), 185–194.

[24] L. Ma, L. Zhao, *Classification of positive solitary solutions of the nonlinear Choquard equation*, Arch. Rational Mech. Anal., 195 (2010), 455–467.

[25] V. Moroz, J. Van Schaftingen, *Nonexistence and optimal decay of supersolutions to Choquard equations in exterior domains*, J. Differential Equations, **254** (2013), 3089–3145.

[26] K. Nakanishi, *Energy scattering for Hartree equations*, Math. Res. Lett., **6** (1999), 107–118.

[27] P. Quittner, P. Souplet, *Symmetry of components for semilinear elliptic systems*, SIAM J. Math. Anal., **44** (2012), 2545–2559.

[28] D. Smets, *Nonlinear Schrödinger equations with Hardy potential and critical nonlinearities*, Trans. Amer. Math. Soc., **357** (2005), 2909–2938.